

Mental Health

**Compendium of Health Related Statistics
April 2005**


Our Healthier Nation

**To reduce the death rate from suicide and undetermined injury
by at least a fifth by 2010**

**Mortality from suicide and injury undetermined (ICD9 E950-E959, E980-E989 exc E988.8; ICD10 X60-X84, Y10-Y34 exc Y33.9)
Directly standardised rates per 100,000 (and 95% confidence intervals), 2001-02 (and 1995-97
baseline years) Pooled, all ages**

		Baseline Rate 1995-97		2001 & 2002				Target Rate 2010
		Rate	No. of deaths	Rate	No. of deaths	Lower Confidence Interval	Upper Confidence Interval	Rate
Gateshead	Males	17.39	52	12.47	22	7.20	17.74	13.91
	Females	5.99	18	3.48	7	0.86	6.09	4.79
	Persons	11.57	70	7.77	29	4.89	10.65	9.26
North East	Males	16.01	614	15.34	382	13.79	16.90	12.81
	Females	4.21	172	3.91	108	3.15	4.66	3.37
	Persons	9.92	786	9.45	490	8.60	10.30	7.94
England	Males	14.1	10312	13.42	6724	13.10	13.74	11.28
	Females	4.47	3563	4.2	2286	4.03	4.38	3.58
	Persons	9.16	13875	8.72	9010	8.54	8.90	7.33
England & Wales	Males	14.24	11023	13.62	7209	13.30	13.93	11.39
	Females	4.45	3753	4.2	2421	4.03	4.38	3.56
	Persons	9.22	14776	8.82	9630	8.64	8.99	7.38

Source: Compendium of Clinical & Health Indicators 2003, ONS




Each individual year shown is based on a three-year period, so that for instance 1998 represents 1997-99.



Our Healthier Nation

Indirectly Age-Standardised Mortality Ratios (SMR and 95% Confidence Intervals), 2001-02
(ICD10 X60-X84):

		Number of deaths	SMR	Lower Confidence Interval	Upper Confidence Interval
Gateshead	Males	12	64	33	113
	Females	5	90	29	210
	Persons	17	70	41	113
North East	Males	266	108	95	122
	Females	67	92	71	116
	Persons	333	104	93	116
England & Wales	Males	5,169	100	97	103
	Females	1,498	100	95	105
	Persons	6,667	100	98	102

Source: Compendium of Clinical and Health Indicators 2003





Source: Compendium of Clinical and Health Indicators 2003


Note:



In 1993 ONS introduced changes in the way cause of death information is coded. As a result, data for 1993 and subsequent years are not comparable with data from preceding years

**Our Healthier Nation
 Purposely Inflicted: All Ages
 Indirectly Age-Standardised Mortality Ratios (SMR and 95% Confidence Intervals), 2001-02
 (ICD10 X60-X84, Y10-Y34 exc. Y33.9)**

		Number of deaths	SMR	Lower Confidence Interval	Upper Confidence Interval
Gateshead	Males	22	85	53	128
	Females	7	78	31	161
	Persons	29	83	56	119
North East	Males	382	111	100	123
	Females	108	92	75	111
	Persons	490	106	97	116
England & Wales	Males	7209	100	98	102
	Females	2421	100	96	104
	Persons	9630	100	98	102

Source: Compendium of Clinical and Health Indicators 2003





Source: Compendium of Clinical and Health Indicators 2003

Note:

In 1993 ONS introduced changes in the way cause of death information is coded. As a result, data for 1993 and subsequent years are not comparable with data for preceding years.

**Hospital Episodes Associated with Schizophrenia (ICD10 F20, F21, F23.2, F25)
Directly Standardised Rates, Ages 15-74, Financial Year 2001/02
(Age-standardised rates per 100,000 population)**

		Number of FCEs	Rate per 100,000	Lower Confidence Interval	Upper
Gateshead	Males	98	149.70	119.78	179.62
	Females	56	77.83	57.28	98.38
	Persons	154	113.69	95.55	131.83
North East	Males	950	105.52	98.77	112.27
	Females	491	50.81	46.28	55.34
	Persons	1441	77.79	73.75	81.83
England	Males	17981	99.22	97.76	100.68
	Females	10028	53.70	52.64	54.76
	Persons	28009	76.46	75.56	77.36

Source: Compendium of Clinical and Health Indicators 2003

Indicator is different to previous indicator of associated with schizophrenia due to there being only a small number of deaths mentioning schizophrenia and they are usually among elderly people who have been in long-stay institutions for many years, not suicides of young people suffering from schizophrenia. In addition, the ICD-9 295 and ICD-10 F20-F29 codes do not appear to have exact equivalents.